 **UDENYCA** **ONBODY**[®]
(pegfilgrastim-cbqv)
Injection 6 mg/0.6 mL

Actor portrayal

On-the-go treatment injected *IN JUST 5 MINUTES*



What is UDENYCA?

UDENYCA (pegfilgrastim-cbqv) is a prescription medicine used to help reduce the chance of infection due to a low white blood cell count, in people with certain types of cancer (non-myeloid), who receive anti-cancer medicines (chemotherapy) that can cause fever and low white blood cell count.

IMPORTANT SAFETY INFORMATION

Do not take UDENYCA if you have had a serious allergic reaction to pegfilgrastim or filgrastim products.

Please see additional Important Safety Information throughout this brochure and full [Prescribing Information](#) and [Patient Information](#).

UNDERGOING CHEMOTHERAPY?


UDENYCA can help support your immune system



Chemotherapy can decrease your white blood cell count

A decrease in your white blood cell count can increase your risk of:

- Infection
- A complication called febrile neutropenia, which can delay your chemotherapy treatment



UDENYCA boosts the growth of your white blood cells, which may help your body:

- **Fight off** infection
- **Prevent** febrile neutropenia

The most common side effects of UDENYCA are pain in the bones, arms, and legs

These are not all of the possible side effects of UDENYCA. Please see the Important Safety Information throughout this brochure and contact your healthcare team for medical advice about side effects or concerns you may have about your treatment.

Together, you can decide on the best way to manage side effects so you can stay on track with your treatment.

How do you know if you have an infection?



A fever is a sign of an infection and a medical emergency

- If you experience a fever of, or higher than, 100.4 °F (38.0 °C), call your healthcare team immediately or go to the hospital
- Take your temperature any time you don't feel well or as recommended by your healthcare team



Other signs of infection may include:

- Chills
- Shortness of breath
- Sore throat
- Stiff neck
- Burning or pain with urination

IMPORTANT SAFETY INFORMATION (continued)

Before taking UDENYCA, tell your healthcare provider about all of your medical conditions, including if you:

- Have a sickle cell disorder
- Have had severe skin reactions to acrylic adhesives
- Have kidney problems
- Are pregnant or plan to become pregnant. It is not known if UDENYCA will harm your unborn baby
- Are breastfeeding or plan to breastfeed. It is not known if UDENYCA passes into your breast milk

Please see additional Important Safety Information throughout this brochure and full [Prescribing Information](#) and [Patient Information](#).



Learn more about your treatment at UDENYCA.com



Actor portrayals

IMPORTANT SAFETY INFORMATION (continued)


Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Know the medicines you take. Keep a list of them to show your healthcare provider and pharmacist when you get a new medicine.

ON THE GO?

The on-body injector delivers your treatment in just 5 minutes*

Black fill indicator shows status of UDENYCA delivery

Status light signals if the on-body injector is working properly



High-quality, well-tolerated adhesive for secure attachment on your body



Preset to **automatically deliver** your dose of UDENYCA in **just 5 minutes***

Status light guide



Flashing green

The on-body injector is working properly. Do not remove the on-body injector if the status light is flashing green.



Solid green (or off)

Signals dose delivery is complete. Check to see if fill indicator reads empty.



Flashing red

On-body injector error. If you hear beeping and see a flashing red light, call your healthcare provider immediately—you may need a replacement dose.

*Please refer to the [Instructions for Use](#) before receiving UDENYCA.

IMPORTANT SAFETY INFORMATION

What are the possible side effects of UDENYCA?

UDENYCA can cause serious side effects, including:

Spleen rupture: Your spleen may become enlarged and can rupture. A ruptured spleen can cause death. Call your healthcare provider right away if you have pain in the left upper stomach (abdomen) area or your left shoulder.

A serious lung problem called acute respiratory distress syndrome (ARDS): Call your healthcare provider or get emergency medical help right away if you have shortness of breath with or without a fever, trouble breathing, or a fast rate of breathing.


Please see additional Important Safety Information throughout this brochure and full [Prescribing Information](#) and [Patient Information](#).






What to expect with your on-body injector

The steps below can help you understand what to expect while wearing the on-body injector, but they do not replace the [Instructions for Use](#). Please refer to it while wearing the on-body injector. Talk to your healthcare provider or pharmacist if you have any questions.



For disposal information, visit [fda.gov/safesharpsdisposal](https://www.fda.gov/safesharpsdisposal).

-  **If you hear beeping at any time, check the status light. If it is flashing red, call your healthcare provider right away.**
- If you have an allergic reaction during the delivery of UDENYCA, remove the on-body injector and call your healthcare provider or seek emergency care right away.**

The UDENYCA on-body injector may be right for you if you:



Actor portrayal



Don't want to return for a next-day office visit



Live far away from your doctor's office or depend on a caregiver for a ride



Prefer your healthcare provider set up your treatment



Have a weakened immune system and need to limit exposure to infection at the doctor's office

IMPORTANT SAFETY INFORMATION

What are the possible side effects of UDENYCA? (continued)

Serious allergic reactions: UDENYCA can cause serious allergic reactions. These reactions can cause a rash over your whole body, shortness of breath, wheezing, dizziness, swelling around your mouth or eyes, fast heart rate, and sweating. **If you have any of these symptoms, stop using UDENYCA and call your healthcare provider or get emergency medical help right away.**

Sickle cell crisis: You may have a serious sickle cell crisis, which could lead to death, if you have a sickle cell disorder and receive UDENYCA. Call your healthcare provider right away if you have symptoms of sickle cell crisis such as pain or difficulty breathing.

Kidney injury (glomerulonephritis): UDENYCA can cause kidney injury. Call your healthcare provider right away if you experience swelling of your face or ankles, blood in your urine (or dark-colored urine), or you urinate less than usual.

Please see additional Important Safety Information throughout this brochure and full [Prescribing Information](#) and [Patient Information](#).



The dos and don'ts of using the on-body injector

WATCH THE VIDEO
to learn what to expect
with the on-body injector



To ensure the proper use of your on-body injector and to achieve the best possible outcomes with UDENYCA, please follow these guidelines:

DO

- ✔ Be careful not to bump the on-body injector or knock the on-body injector off your body.
- ✔ **WEAR THE ON-BODY INJECTOR UNDER CLOTHING.** Ensure the on-body injector is protected from direct sunlight. If it is exposed to sunlight for more than 1 hour, it may affect your medication.
- ✔ **CHECK THE STATUS LIGHT REGULARLY.**
- ✔ **CHECK YOUR ON-BODY INJECTOR OFTEN FOR LEAKS** during the 5-minute dose delivery. After 5 minutes, check to see if your on-body injector is empty.
- ✔ **AFTER SHOWERING,** check the on-body injector to make sure it has not become loose or dislodged.

DON'T

- ✘ **DO NOT PEEL OFF OR DISTURB THE ON-BODY INJECTOR ADHESIVE** before your full dose is complete. Interrupting the on-body injector before the dose is complete may result in a missed or incomplete dose of UDENYCA.
- ✘ **DO NOT REMOVE THE ON-BODY INJECTOR IF THE STATUS LIGHT IS FLASHING GREEN.** The on-body injector is still working properly, and your dose is not yet complete.
- ✘ **DO NOT EXPOSE** to diagnostic imaging (eg, CT scan, MRI, Ultrasound, X-ray), radiation treatment, or oxygen-rich environments (such as hyperbaric chambers).
- ✘ **DO NOT ALLOW LOTIONS, CREAMS, OILS, OR CLEANING AGENTS** to come into contact with the on-body injector at any time. The day before your next UDENYCA dose, avoid using lotions, creams, and oils on the area where the injector will be attached.
- ✘ **DO NOT USE** bathtubs, hot tubs, whirlpools, or saunas while wearing the on-body injector. This may affect your medicine and delivery.
- ✘ **DO NOT SLEEP ON OR APPLY PRESSURE** to the on-body injector during wear, especially during dose delivery. This may affect the on-body injector performance.

Please refer to the [Patient Instructions for Use](#) for additional details regarding dos and don'ts when wearing your on-body injector.





To be completed by your healthcare provider after you receive the UDENYCA on-body injector

On-body injector was applied at:

Injection of your dose will start around:

If you have questions, call us at:

IMPORTANT SAFETY INFORMATION

What are the possible side effects of UDENYCA? (continued)

Increased white blood cell count (leukocytosis): Your healthcare provider will check your blood during treatment with UDENYCA.

Decreased platelet count (thrombocytopenia): Your healthcare provider will check your blood during treatment with UDENYCA. Tell your healthcare provider if you have unusual bleeding or bruising during treatment with UDENYCA. This could be a sign of decreased platelet counts, which may reduce your blood's ability to clot.

Capillary leak syndrome: Pegfilgrastim products can cause fluid to leak from blood vessels into your body's tissues. This condition is called "Capillary Leak Syndrome" (CLS). CLS can quickly cause you to have symptoms that may become life-threatening. Get emergency medical help right away if you develop any of the following symptoms:

- Swelling or puffiness and urinating less than usual
- Trouble breathing
- Swelling of your stomach area (abdomen) and feeling of fullness
- Dizziness or feeling faint
- A general feeling of tiredness

Myelodysplastic syndrome (MDS) and acute myeloid leukemia (AML): If you have breast cancer or lung cancer and are receiving UDENYCA in combination with chemotherapy and radiation therapy, or radiation therapy alone, you may have an increased risk of developing a precancerous blood condition called MDS or a blood cancer called AML. Symptoms of MDS and AML may include tiredness, fever, and easy bruising or bleeding. Call your healthcare provider if you develop these symptoms during treatment with UDENYCA.



Financial support is available

AccordCares® provides financial support options for eligible patients prescribed UDENYCA. Ask your doctor's office for help applying.



FIND MORE DETAILS

about support and eligibility by visiting [UDENYCA.com/AccordCares](https://www.udenyca.com/accordcares), or call [1-844-483-3692](tel:1-844-483-3692).

Inflammation of the aorta (aortitis): Inflammation of the aorta (the large blood vessel that transports blood from the heart to the body) has been reported in patients who received pegfilgrastim products. Symptoms may include fever, abdominal pain, feeling tired, and back pain. Call your healthcare provider if you experience these symptoms.

Allergies to acrylics (UDENYCA ONBODY® only): The on-body injector (OBI) for UDENYCA uses acrylic adhesive. If you have had reactions to acrylic adhesives, use of this product may result in a significant reaction.

The most common side effects of UDENYCA are pain in the bones, arms, and legs.

These are not all the possible side effects of UDENYCA. Tell your healthcare provider about any side effect you experience. For additional information, see the full Prescribing Information or ask your healthcare provider or pharmacist.

You are encouraged to report side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call [1-800-FDA-1088](tel:1-800-FDA-1088).

UDENYCA is supplied as:

- UDENYCA ONBODY®: 6 mg/0.6 mL in a single-dose prefilled syringe co-packaged with the on-body injector for UDENYCA®.

Please see additional Important Safety Information throughout this brochure and full [Prescribing Information](#) and [Patient Information](#).



Accord, AccordCares, UDENYCA ONBODY, UDENYCA, and associated logos are trademarks of Accord BioPharma, Inc. and its licensors or affiliates.

©2026 Accord BioPharma. All rights reserved. This content is intended for U.S. audiences only. US-UDY-2500010 01/26