

Reduce your risk of chemotherapy-related infection.
Choose from 3 administration options to fit your needs.

Immune system support that *FITS INTO YOUR LIFE*



ON THE GO

AT HOME

IN THE OFFICE

Actor portrayals

What is UDENYCA?

UDENYCA (pegfilgrastim-cbqv) is a prescription medicine used to help reduce the chance of infection due to a low white blood cell count, in people with certain types of cancer (non-myeloid), who receive anti-cancer medicines (chemotherapy) that can cause fever and low white blood cell count.

IMPORTANT SAFETY INFORMATION

Do not take UDENYCA if you have had a serious allergic reaction to pegfilgrastim or filgrastim products.

Before taking UDENYCA, tell your healthcare provider about all of your medical conditions, including if you:

- Have a sickle cell disorder
- Have had severe skin reactions to acrylic adhesives
- Have kidney problems
- Are pregnant or plan to become pregnant. It is not known if UDENYCA will harm your unborn baby
- Are breastfeeding or plan to breastfeed. It is not known if UDENYCA passes into your breast milk

Please see additional Important Safety Information throughout this brochure and full [Prescribing Information](#).


UNDERGOING CHEMOTHERAPY?

UDENYCA can help support your immune system

Chemotherapy can decrease your white blood cell count

A decrease in your white blood cell count can increase your risk of:

- Infection
- A complication called febrile neutropenia, which can delay your chemotherapy treatment



UDENYCA boosts the growth of your white blood cells, which may help your body:

- **Fight off** infection
- **Prevent** febrile neutropenia

The most common side effects of UDENYCA are pain in the bones, arms, and legs

These are not all of the possible side effects of UDENYCA. Please see the Important Safety Information throughout this brochure and contact your healthcare team for medical advice about side effects or concerns you may have about your treatment.

Together, you can decide on the best way to manage side effects so you can stay on track with your treatment.

How do you know if you have an infection?



A fever is a sign of an infection and a medical emergency

- If you experience a fever of, or higher than, **100.4 °F (38.0 °C)**, call your healthcare team immediately or go to the hospital
- Take your temperature any time you don't feel well or as recommended by your healthcare team



Other signs of infection may include:

- Chills
- Shortness of breath
- Sore throat
- Stiff neck
- Burning or pain with urination

IMPORTANT SAFETY INFORMATION (continued)

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Know the medicines you take. Keep a list of them to show your healthcare provider and pharmacist when you get a new medicine.



Learn more about your
treatment at [UDENYCA.com](https://www.udenyca.com)



Actor portrayals

IMPORTANT SAFETY INFORMATION (continued)

What are the possible side effects of UDENYCA?

UDENYCA can cause serious side effects, including:

Spleen rupture: Your spleen may become enlarged and can rupture. A ruptured spleen can cause death. Call your healthcare provider right away if you have pain in the left upper stomach (abdomen) area or your left shoulder.

A serious lung problem called acute respiratory distress syndrome (ARDS): Call your healthcare provider or get emergency medical help right away if you have shortness of breath with or without a fever, trouble breathing, or a fast rate of breathing.

ON THE GO?


The on-body injector delivers your treatment in just 5 minutes*

High-quality, well-tolerated adhesive for secure attachment on your body

Black fill indicator

shows status of UDENYCA delivery

Status light signals if the on-body injector is working properly



Preset to **automatically deliver** your dose of UDENYCA in **just 5 minutes***

Status light guide



Flashing green

The on-body injector is working properly. **Do not** remove the on-body injector if the status light is flashing green.



Solid green (or off)

Signals dose delivery is complete. Check to see if fill indicator reads empty.



Flashing red

On-body injector error. If you hear beeping and see a flashing red light, call your healthcare provider immediately—you may need a replacement dose.

*Please refer to the [Instructions for Use](#) before receiving UDENYCA.

IMPORTANT SAFETY INFORMATION (continued)

What are the possible side effects of UDENYCA? (continued)

Serious allergic reactions: UDENYCA can cause serious allergic reactions. These reactions can cause a rash over your whole body, shortness of breath, wheezing, dizziness, swelling around your mouth or eyes, fast heart rate, and sweating. **If you have any of these symptoms, stop using UDENYCA and call your healthcare provider or get emergency medical help right away.**

Sickle cell crisis: You may have a serious sickle cell crisis, which could lead to death, if you have a sickle cell disorder and receive UDENYCA. Call your healthcare provider right away if you have symptoms of sickle cell crisis such as pain or difficulty breathing.

WATCH THE VIDEO
 Learn what to expect with the
 UDENYCA on-body injector



What to expect with your on-body injector

The steps below can help you understand what to expect while wearing the on-body injector, but they do not replace the **Instructions for Use**. Please refer to it while wearing the on-body injector. Talk to your healthcare provider or pharmacist if you have any questions.

Begin




Day you receive chemotherapy

Your healthcare provider will apply your on-body injector. The status light will flash green, indicating that it is working properly.

Note: After it's applied, **keep it dry for the next 3 hours**. This is to ensure proper adhesion and functionality of the device.


Hours 1 to 27



Monitor your on-body injector

Occasionally check the status light to **confirm it's flashing green**. If applied on the back of your arm, a caregiver must be available to confirm. **Ensure that on-body injector is dry for 3 hours before dose delivery.**


Hour 27



Injection begins

The on-body injector needle will automatically insert into your skin. During the injection, the on-body injector will flash a fast, green light, and you may hear the pump working. It will take just **5 minutes** for your dose of UDENYCA to be delivered.


**Hour 27
 + 5 minutes**



Injection is complete

When the injection is complete, a long beep will sound, and the status light will turn solid green. Confirm your injection is complete by checking that the status light is solid green or has turned off.


Complete



Remove the on-body injector

Grasp the edge of the adhesive and slowly peel off the device. Do not grab the on-body injector to pull it off. Check to see if your on-body injector is empty. You should see a black line next to the **EMPTY** indicator.

For disposal information, go to page 9 or visit fda.gov/safesharpsdisposal.

-  If you hear beeping at any time, check the status light. If it is flashing red, call your healthcare provider right away.
- If you have an allergic reaction during the delivery of UDENYCA, remove the on-body injector and call your healthcare provider or seek emergency care right away.

WANT TREATMENT FROM THE COMFORT OF HOME?*

The autoinjector can be administered by you or a care partner

Gray needle guard covers the needle and **keeps you safe** from needlesticks

Clear cap covers the gray needle guard

Large, clear viewing window allows you to **see when your medicine has been injected**. An orange indicator fills the window when your injection is complete.



Delivers medicine in **less than 10 seconds**[†]



Label with expiration date

*May also be given by your provider in their office.

[†]Please refer to the [Instructions for Use](#) before administering UDENYCA.

IMPORTANT SAFETY INFORMATION (continued)

What are the possible side effects of UDENYCA? (continued)

Kidney injury (glomerulonephritis): UDENYCA can cause kidney injury. Call your healthcare provider right away if you experience swelling of your face or ankles, blood in your urine (or dark-colored urine), or you urinate less than usual.

Increased white blood cell count (leukocytosis): Your healthcare provider will check your blood during treatment with UDENYCA.

Decreased platelet count (thrombocytopenia): Your healthcare provider will check your blood during treatment with UDENYCA. Tell your healthcare provider if you have unusual bleeding or bruising during treatment with UDENYCA. This could be a sign of decreased platelet counts, which may reduce your blood's ability to clot.

WATCH THE VIDEO

See step-by-step instructions on using the UDENYCA autoinjector



A quick guide to the UDENYCA autoinjector

Your healthcare provider will show you how to prepare and inject the UDENYCA autoinjector. This guide does not replace the [Instructions for Use](#). Please refer to it before administering. If you have questions, talk to your healthcare team or pharmacist.

Only adults can self-inject with the UDENYCA autoinjector; keep out of reach of children.

To get started

Have these supplies ready:

- 1 UDENYCA autoinjector
- 1 alcohol wipe
- 1 cotton ball or gauze pad
- 1 FDA-approved sharps container. Learn more on page 9 or at [FDA.gov/safesharpsdisposal](https://www.fda.gov/safesharpsdisposal)

Take the UDENYCA autoinjector out of the refrigerator and bring to room temperature 30 minutes before use.

- Do not warm the autoinjector in any other way. For example, do not warm in a microwave, in hot water, or in direct sunlight
- Dispose of autoinjectors that have been left out at room temperature (68 °F-77 °F) for more than 48 hours


Inspect your autoinjector and do not use if:

- The medicine appears discolored or cloudy
- The medicine contains lumps, flakes, or particles
- It has been dropped, crushed, or damaged in any way
- It has expired



Injecting UDENYCA

1



Choose an injection site.

Clean and dry your hands. Clean the injection site with an alcohol wipe. Stay 2 inches away from the navel if injecting into the abdomen.

2



Remove the clear cap.

Pull the cap straight up. After removing the cap, **do not place it back on the autoinjector**. Do not touch the gray needle guard.

3



Press the autoinjector.

Place the gray needle guard against your skin at a 90° angle. Press the autoinjector against the skin and you will hear the **first click**.

4



Hold the autoinjector

Keep holding the autoinjector firmly against your skin until the orange indicator stops moving and you hear the second click. **After you hear the second click, continue to hold the autoinjector in place and count to 5.**

PREFER IN-OFFICE TREATMENT?

The prefilled syringe is administered by your provider



More than
300,000
patients have
been treated*

Needle guard covers the
needle and keeps you
safe from needlesticks



Latex-free needle cap
may reduce risk in people
with latex sensitivity

*January 2019 through August 2025.

DISCOVER MORE
Learn about the UDENYCA
prefilled syringe



IMPORTANT SAFETY INFORMATION (continued)

What are the possible side effects of UDENYCA? (continued)

Capillary leak syndrome: Pegfilgrastim products can cause fluid to leak from blood vessels into your body's tissues. This condition is called "Capillary Leak Syndrome" (CLS). CLS can quickly cause you to have symptoms that may become life-threatening. Get emergency medical help right away if you develop any of the following symptoms:

- Swelling or puffiness and urinating less than usual
- Trouble breathing
- Swelling of your stomach area (abdomen) and feeling of fullness
- Dizziness or feeling faint
- A general feeling of tiredness

How to dispose of your used UDENYCA on-body injector or autoinjector*



Put your used UDENYCA product in an FDA-cleared sharps disposal container immediately after use. Do not throw it away in the household trash.

If you do not have an FDA-cleared sharps disposal container, you may use a household container that:

- Is made of a heavy-duty plastic
- Can be closed with a tight-fitting, puncture-resistant lid that prevents sharps from falling out
- Is upright and stable during use
- Is leak-resistant
- Is properly labeled to warn of hazardous waste inside the container



For more information about safe sharps disposal, and for specific information about sharps disposal in the state that you live in, go to [fda.gov/safesharpsdisposal](https://www.fda.gov/safesharpsdisposal).

*The UDENYCA prefilled syringe may also be administered at home. Follow the same sharps disposal steps.

IMPORTANT SAFETY INFORMATION (continued)

What are the possible side effects of UDENYCA? (continued)

Myelodysplastic syndrome (MDS) and acute myeloid leukemia (AML): If you have breast cancer or lung cancer and are receiving UDENYCA in combination with chemotherapy and radiation therapy, or radiation therapy alone, you may have an increased risk of developing a precancerous blood condition called MDS or a blood cancer called AML. Symptoms of MDS and AML may include tiredness, fever, and easy bruising or bleeding. Call your healthcare provider if you develop these symptoms during treatment with UDENYCA.



Financial resources are available

Co-Pay Program

AccordCares offers a Co-Pay Program for eligible patients with commercial insurance that covers out-of-pocket expenses related to UDENYCA.

- Eligible patients may pay as little as **\$0 co-pay** per UDENYCA treatment
- Annual maximum benefit of \$15,000 for out-of-pocket expenses for UDENYCA, including co-pays or co-insurance

\$0 co-pay per UDENYCA treatment*

*For eligible patients.



Independent foundation support

AccordCares may be able to help you find financial support through charitable foundations, or you can choose to contact independent assistance foundations directly.

Ask your doctor's office to help you apply for UDENYCA financial support

Get more details about support and eligibility at UDENYCA.com/AccordCares, or call 1-844-483-3692.



IMPORTANT SAFETY INFORMATION (continued)

What are the possible side effects of UDENYCA? (continued)

Inflammation of the aorta (aortitis): Inflammation of the aorta (the large blood vessel that transports blood from the heart to the body) has been reported in patients who received pegfilgrastim products. Symptoms may include fever, abdominal pain, feeling tired, and back pain. Call your healthcare provider if you experience these symptoms.

Allergies to acrylics (UDENYCA ONBODY® only): The on-body injector (OBI) for UDENYCA uses acrylic adhesive. If you have had reactions to acrylic adhesives, use of this product may result in a significant reaction.



Treatment your way—with UDENYCA



PREFILLED AUTOINJECTOR

Want treatment from the comfort of home?

The autoinjector can be administered by you or a care partner

ON-BODY INJECTOR

On the go?

The on-body injector delivers your treatment in just 5 minutes

PREFILLED SYRINGE

Prefer in-office treatment?

The prefilled syringe is administered by your healthcare provider

IMPORTANT SAFETY INFORMATION (continued)

What are the possible side effects of UDENYCA? (continued)

The most common side effects of UDENYCA are pain in the bones, arms, and legs.

These are not all the possible side effects of UDENYCA. Tell your healthcare provider about any side effect you experience. For additional information, see the full [Prescribing Information](#) or ask your healthcare provider or pharmacist.

You are encouraged to report side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

Please see additional Important Safety Information throughout this brochure and full [Prescribing Information](#).



GET TO KNOW
UDENYCA